arXiv:1203.6419v2 [quant-ph] 9 Apr 2012

From blockade to transparency: controllable photon transmission through a circuit QED system

Yu-xi Liu,[1,2,3] Xun-Wei Xu,[1] Adam Miranowicz,[4,3] and Franco Nori[3,5]
[1]*Institute of Microelectronics, Tsinghua University, Beijing 100084, China*
[2]*Tsinghua National Laboratory for Information Science and Technology (TNList), Tsinghua University, Beijing 100084, China*
[3]*Advanced Science Institute, RIKEN, Wako-shi, Saitama 351-0198, Japan*
[4]*Faculty of Physics, Adam Mickiewicz University, 61-614 Poznań, Poland*
[5]*Physics Department, The University of Michigan, Ann Arbor, Michigan 48109-1040, USA*
(Dated: March 1, 2013)

A strong photon-photon nonlinear interaction is a necessary condition for photon blockade. Moreover, this nonlinearity can also result a bistable behavior in the cavity field. We analyze the relation between detecting field and photon blockade in a superconducting circuit QED system, and show that the photon blockade cannot occur when the detecting field is in the bistable regime. We further demonstrate that the photon transmission through such system can be controlled (from photon blockade to transparency) by the detecting field. Numerical simulations show that our proposal is experimentally realizable with current technology.

PACS numbers: 42.50.Gy, 42.65.Pc, 42.50.Ar, 85.25.-j

Introduction.—Superconducting circuit quantum electrodynamics (QED) allows studying the interaction between artificial superconducting atoms and quantized microwave fields (see, e.g., reviews [1, 2] and references therein). The coupling strengths have been explored from the strong regime to the ultrastrong one [3–6]. It is well known that the equally-spaced energy structure of the quantized microwave field can be changed to an anharmonic one, including dressed states [7], formed by a superconducting charge qubit and classical microwave field, which have been observed experimentally [8].

The nonlinear energy splitting in circuit QED has been shown by experimentally analyzing the Rabi frequencies for different numbers of microwave quanta inside the cavity [9]. Moreover, the experimental spectrum for the square root of the photon-number nonlinearity was reported [10]. Furthermore, the nonlinear response of the vacuum Rabi resonance was demonstrated [11]. Current experimental data indicate that the nonlinearity of the microwave photons can be many orders of magnitude larger than that of macroscopic media. All of these experiments [8–11] lay a solid foundation for the development of microwave nonlinear interactions on a chip, and open the door to further study microwave nonlinear quantum optics at the level of single artificial atoms and single microwave photons. For example, photon blockade [12], which is the analog of the Coulomb blockade for electrons in single-electron devices, has recently been observed in: an optical cavity with one trapped atom [13], quantum dots coupled to the photonic crystal cavity [14], and also superconducting circuit QED with resonant [15] and dispersive [16] atom-field interactions.

Photon blockade originates from the photon-photon interaction, which prevents subsequent photons from resonantly entering a cavity. Therefore, photon blockade can be used as a single-photon source or single-photon transistor. The microwave nonlinear interaction might also be used to improve superconducting qubit readout [17, 18]. One of the basic conditions for photon blockade is that the decay rate of the cavity field should be less than the photon-photon interacting strength. We know that the decay rates of the photon number states are very different from those of coherent states [19, 20], thus the blockaded photons might not be observed when the photon number inside the cavity is extremely large. In addition, bistability is one of basic properties of nonlinear systems, but there is lack of studies of bistability on the photon blockade.



FIG. 1: (Color online) (a) Eigenvalues $E_k(n_c)$ (green curves) of the free Hamiltonian $\hbar\kappa(n-n_c)$ versus n_c for the photon numbers $n=0$ and $n=1$ inside the cavity, in energy units of $\hbar\kappa$. Blue curves are the eigenvalues near $n_c=0.5$, when the external field is included. The degeneracy of the two eigenvalues in the free Hamiltonian is lifted at the point $n_c=0.5$, with distance $\hbar|\Omega|$ by the external field. (b) Several eigenvalues $E_k(n_c)$ (up to four) of the Hamiltonian in Eq. (2) vary with n_c for the ratio $\kappa/|\Omega|=10$. (c) The mean photon number in the ground state varies with n_c for $\kappa/|\Omega|=1$ (green dash-dotted curve), 10 (red dashed curve), and 100 (blue solid curve). This is the photon analog of the Coulomb staircase.

Here we focus on the Jaynes-Cummings model of circuit QED in the strong-coupling regime, study the relation between bistability and photon blockade, and analyze the effect of the detecting field on the photon blockade by studying the second-order degrees of coherence for the cavity field, explore the possibility of unblocking the photon of the weak detecting (or signal) field by using electromagnetically-induced transparency [22].

Model.—We start from the standard Jaynes-Cummings model for circuit QED with a transmission line resonator coupled to a superconducting qubit with coupling strength g. The

TABLE I: Equivalence between photon and Coulomb blockades (or similarity between photon blockade and superconducting charge qubits).

Quanta	Photons	Electrons
Characteristic energy	Kerr energy $\hbar\kappa$	Charging energy E_c
Energy offset control	Driving field frequency ω_d	Gate voltage V_g
Detecting method	Driving field $E(t)$	Bias voltage V_b
Measurement of output	Photon number $\langle n\rangle$	Electron current I
Tunneling energy	Coupling energy $\hbar\|\Omega\|$	Josephson energy E_J

frequencies of the cavity field and the qubit are ω_0 and ω_q, respectively. If the cavity is driven by classical fields, then with a canonical transformation [21], the driven Hamiltonian is transformed to the dressed one

$$H = \hbar\omega_0 a^\dagger a + \frac{\hbar}{2}\left[\omega_0 - E(N)\right]\sigma_z + \hbar\left[E(t)a^\dagger + E^*(t)a\right], \tag{1}$$

with the creation $a^\dagger$ (annihilation a) operator of the cavity field and $E(N) = \sqrt{\Delta^2 + 4g^2N}$. Here the operator $N = a^\dagger a + (\sigma_z + 1)/2$ describes the total number of excitations for the photon and the qubit, and $\Delta = \omega_0 - \omega_q$ describes the detuning between the cavity field and the qubit. The third term in the right of Eq. (1), with the parameter $E(t)$, describes the interaction between the external fields and the cavity mode. In the derivation of Eq. (1), the terms proportion to $O(N^{-1/2})$ are neglected. The excited and ground states with sign ± for σ_z are considered as the attractive and repulsive photon-photon interactions. Below we assume that the qubit is always in its excited state. We also assume that the decay of the qubit is negligibly small in the bad-cavity limit.

If a monochromatic driving field (or saying the detecting field for photon blockade), with frequency ω_d, is coupled to the cavity via $E(t) = \Omega e^{-i\omega_d t}$ with coupling constant Ω, then the Hamiltonian in Eq. (1) can be used to describe photon blockade [12] when the amplitude of Ω for the driving field is much smaller than the photon-photon coupling constant κ given below. Moreover the decay rate γ of the cavity field should also be much smaller than κ to experimentally observe photon blockade. In the dispersive regime, the single two-level atom-induced photon blockade can be understood by expanding the Hamiltonian in Eq. (1) up to third order in parameter g/Δ as

$$H_{\rm eff} = \hbar\kappa(n - n_{\rm c})^2 + \hbar(\Omega a^\dagger + \Omega^* a) \tag{2}$$

with $n = a^\dagger a$ in the rotating reference frame with frequency ω_d for the driving field. Here, $\kappa = g^4/\Delta^3$ denotes the photon-photon interaction strength, $n_c = (\kappa - \Delta_d)/2\kappa$ is the rescaled detuning $\Delta_d = \omega_0 - \omega_d$ between the driving and the cavity fields. In the derivation of Eq. (2), several constant terms and also the qubit state-dependent cavity-frequency shift have been neglected with the assumption that the qubit is in its excited state. Equation (2) shows that $\hbar\kappa$ and ω_d (or $n_{\rm c}$) are equivalent to the charging energy and gate voltage in single-electron devices for Coulomb blockade. Equation (2) also shows that the parameters κ, ω_d (or $n_{\rm c}$), and $|\Omega|$ play similar functions as the charging energy, the gate voltage, and the Josephson energy of the charge qubit (see, e.g., Ref. [23]). For resonant driving, $\Delta_d = 0$ and $n_{\rm c} = 0.5$, the photon states $|0\rangle$ and $|1\rangle$ are degenerate for the free Hamiltonian $\hbar\kappa(n - 0.5)^2$. As shown in Fig. 1(a), this degeneracy can be lifted by the coupling strength $|\Omega|$. Figures 1(b,c) show the variations of the eigenenergies and the mean photon number $\langle n\rangle$ of the ground state for the Hamiltonian in Eq. (2) as functions of the parameter $n_{\rm c}$. Variations for both the eigenenergy and the mean photon number are the same as those of the eigenenergy and mean charge number of charge qubits or single-electron devices for Coulomb blockade. The staircase shape for the mean photon number was experimentally demonstrated in circuit QED system [16]. Moreover, Fig. 1(a) also shows that a large ratio $\kappa/|\Omega|$ corresponds to a sharper step, i.e., a weak driving field and strong photon-photon interaction are more useful for detecting photon blockade. Photon and Coulomb blockades are compared in Table I.



FIG. 2: (Color online) Logarithm $\log(|A_s|)$ of the steady-state solution $|A_s|$ versus the strength of the driving field $|\Omega|$, for different detunings in (a) and (b). Here, $N_{\rm up} = g^4/\Delta^3\gamma = \kappa/\gamma$ is the upper bound photon number for the photon blockade, $\Delta/2\pi = 1$ GHz, $\gamma/2\pi = 0.1$ MHz, and $g/2\pi = 200$ MHz. For above given parameters, $|A_s|$ has two stable states when $0.0091\,\mathrm{MHz} < \Delta_{\rm d} < 36.95\,\mathrm{MHz}$.

Bistability and blockade.—As shown in Fig. 1(c), the basic feature of the staircase shape for the photon blockade gradually disappears when the strength of driving field approaches or is larger than the photon-photon coupling strength. It is known that the driven nonlinear photonic systems exhibit bistability in a regime of parameters. To study the relation between bistability and photon blockade, let us now write the following Heisenberg-Langevin equation

$$\frac{\partial a}{\partial t} = -\left[i\omega_0 + \gamma - i\frac{g^2\sigma_z}{E(N)}\right]a - i\Omega e^{-i\omega_d t} - \sqrt{2\gamma}a_{\rm in}(t) \tag{3}$$

for the driven Hamiltonian in Eq. (1) by the monochromatic classical field $E(t) = \Omega e^{-i\omega_d t}$. The parameter γ is the decay rate of the cavity field. The quantum fluctuations $a_{\rm in}$ due to the vacuum field are assumed to have a zero mean value $\langle a_{\rm in}(t)\rangle = 0$ and Markov correlation $\langle a_{\rm in}(t)a_{\rm in}^\dagger(t')\rangle = \delta(t - t')$. If we denote $a(t) = A(t)e^{-i\omega_d t}$, then the steady-state solution $A_s \equiv \langle A\rangle_s$ for the cavity field becomes

$$A_s = -i\Omega/\{\gamma + i[\Delta_d - g^2\sigma_z/E(\overline{N})]\}, \tag{4}$$



FIG. 3: (Color online) $g^{(2)}(0)$ versus the driving field strength $|\Omega|$ in (a) and $g^{(2)}(\tau)$ versus the delay time τ in (b). Here the other parameters are taken as $\Delta_d = 38.5$ MHz, $\Delta/2\pi = 1$ GHz, $g/2\pi = 200$ MHz, and $\gamma/2\pi = 0.1$ MHz. $g^{(2)}(0)$ versus the detuning Δ_d in (c) for $g/2\pi = 100$ MHz and (d) for $g/2\pi = 200$ MHz; with $\Delta/2\pi = 1$ GHz, $|\Omega|/2\pi = 0.01$ MHz, and $\gamma/2\pi = 0.1$ MHz. For parameters given in (c) and (d), the condition of stable states for the driving field is $0.018\,\mathrm{MHz} < \Delta_\mathrm{d} < 9.627\,\mathrm{MHz}$ and $0.0091\,\mathrm{MHz} < \Delta_\mathrm{d} < 36.95\,\mathrm{MHz}$, respectively.

by using Eq. (3) and the mean-field approximation, e.g., $\langle a^\dagger a\rangle_s = \langle a^\dagger\rangle_s\langle a\rangle_s$. Here $E(\overline{N})$ is the expression of $E(N)$ with replacing operator $a^\dagger a$ in N by steady state value $|A_s|^2$.

Equation (4) shows that the cavity field amplitude $|A_s|^2$ has two stable states when Δ_d satisfies the condition $\Delta_- < \Delta_d < \Delta_+$ with $\Delta_\pm = [g^2/E(\overline{N})] - \{2g^4|A_s|^2 \mp [4g^8|A_s|^4 - \gamma^2E^6(\overline{N})]^{1/2}\}/E^3(\overline{N})]$. Equation (4) also clearly shows when the qubit is decoupled from the cavity field (i.e., $g = 0$) or when the driving field makes $|A_s|^2$ extremely large, such that $g\sigma_z/E(\overline{N}) \approx 0$, then the response of the system is the same as that of the harmonic oscillator. However when $g \neq 0$, the resonant peak of $|A_s|^2$ will move to $\omega_d = \omega_0 - g\sigma_z/E(\overline{N})$. That is, when the cavity contains $|A_s|^2$ photons, the frequency ω_d of the driving field should be increased an amount $-g\sigma_z/E(\overline{N})$ to overcome the photon blockade [12]. The upper bound photon number inside the cavity for the photon blockade is $N_\mathrm{up} \sim g^4/(\Delta^3\gamma) = \kappa/\gamma$ in the dispersive regime as discussed in Ref. [27].

In Fig. 2, the steady state $|A_s|$ versus the input $|\Omega|$ is plotted for several different detunings Δ_d. Figure 2 clearly shows that the bistability disappears for either $\Delta_d > \Delta_+$ or $\Delta_d < \Delta_-$. Figure 2 also shows that most values for $|A_s|$ are smaller than the upper bound value $\sqrt{N_\mathrm{up}}$ for some values of Δ_d in the bistable regime, e.g., values in the region near $\Delta_d/2\pi = 37$ MHz, as shown in Fig. 2(b). However, as shown in Fig. 2 (a) for other parameter regimes of Δ_d, we find that the upper bound value $\sqrt{N_\mathrm{up}}$ can be smaller than some values of $|A_s|$ corresponding to the lower branch of the bistable curve. Therefore, Fig. 2 tells us that N_up is a necessary but not sufficient condition for photon blockade. Because one input corresponds to two stable outputs in the hysteresis region, thus the photon blockade is not well defined in such region.

Dependence of photon blockade on the detuning Δ_d and coupling strength Ω.— To show the effect of the driving field on the photon blockade, let us now study the statistical properties of the cavity field when the input a_in of the vacuum fields in Eq. (3) is considered. We assume that the vacuum fields a_in in Eq. (3) result in a small fluctuation $A_f(t)$ of the cavity field near its stable steady-state A_s by writing the cavity operator as $A(t) = A_s + A_f(t)$. Besides the steady-state solution as in Eq. (4), with the input $a_\mathrm{in}(t) = A_\mathrm{in}(t)e^{-i\omega_d t}$, we can obtain an equation of motion for the fluctuation operator $A_f(t)$ as

$$\frac{\partial A_f(t)}{\partial t} = -\left[i\widetilde{\Delta}_d(\overline{N}) + \gamma\right]A_f(t) - i\alpha(\overline{N})A_f^\dagger(t) - \sqrt{2\gamma}A_\mathrm{in}(t) \tag{5}$$

with $\widetilde{\Delta}_d(\overline{N}) = \Delta_d - \delta\omega(\overline{N})$, $\alpha(\overline{N}) = 2g^4A_s^2\sigma_z/E^3(\overline{N})$, and $\delta\omega(\overline{N}) = g^2\sigma_z[E^2(\overline{N}) - 2g^2|A_s|^2]/E^3(\overline{N})$. Where the terms with higher orders of $A_f(t)$ and $A^\dagger(t)$, e.g., the term $A_f(t)A_f(t)$, have been neglected when Eq. (5) is derived. By applying the Fourier transform $A_f(t) = \int_{-\infty}^{\infty} e^{-i\omega t}A_f(\omega)d\omega/\sqrt{2\pi}$ and also using the conjugate of Eq. (5) with $A_f^\dagger(\omega) = [A_f(-\omega)]^\dagger$, we can easily obtain the solution of the fluctuation as

$$A_f(\omega) = \frac{i\sqrt{2\gamma}}{d(\omega)}\left[\left(\widetilde{\Delta}_\mathrm{d} - \omega + i\gamma\right)A_\mathrm{in}(\omega) + \alpha(\overline{N})A_\mathrm{in}^\dagger(\omega)\right] \tag{6}$$

with the denominator factor $d(\omega) = (i\omega+\gamma)^2 + \widetilde{\Delta}_d^2 - |\alpha(\overline{N})|^2$. The statistical properties of the cavity field can be described via the second-order degrees of coherence $g^{(2)}(\tau)$. Because we assume that the vacuum input is Gaussian and satisfies the Markov correlation, thus $g^{(2)}(\tau)$ can be obtained by just calculating the correlation function with two operators using the Wick's theorem, that is

$$\begin{aligned} g^{(2)}(\tau) &= \frac{2\mathrm{Re}[A_s^2\langle A_f^\dagger(t)A_f^\dagger(t')\rangle + |A_s|^2\langle A_f^\dagger(t)A_f(t')\rangle]}{(|A_s|^2 + \langle n_f(t)\rangle)^2} \\ &+ \frac{|\langle A_f^\dagger(t')A_f^\dagger(t)\rangle|^2 + |\langle A_f^\dagger(t')A_f(t)\rangle|^2}{(|A_s|^2 + \langle n_f(t)\rangle)^2} + 1 \end{aligned} \tag{7}$$

with $t' = t + \tau$ and $\langle n_f(t)\rangle = \langle A_f^\dagger(t)A_f(t)\rangle$. It is easy to obtain $g^{(2)}(\tau)$ straightforwardly by calculating all correlation functions in Eq. (7) using Eq. (6) and $[A_f(-\omega)]^\dagger$. The second-order degrees of coherence $g^{(2)}(0)$ and $g^{(2)}(\tau)$ are plotted in Figs. 3(a,b) using experimentally-accessible parameters. Figures 3(a,b) show that the cavity field tends to the classical behavior when increasing the strength Ω. In this case, the photons might not be blockaded and can transparently pass through the circuit QED system. To explore the effect of the frequency of the driving field, $g^{(2)}(0)$ is plotted as a function of the detuning Δ_d in Figs. 3(c,d) for different coupling strengths between the qubit and the cavity field with other parameters given in the caption of Fig. 3. Figures 3(c,d) clearly demonstrate that the nonclassical behavior of the cavity field is out of the bistable regime for detuning Δ_d. For example, Fig. 3(c) shows $g^{(2)}(0) \leq 1$ when $\Delta_d/2\pi \geq 9.85$ MHz, which is larger than the upper bound value 9.627 Mhz of Δ_d for the bistabilility, and thus the photon blockade cannot occur



FIG. 4: (Color online) The real $A_{\rm R}(\omega')$ and imaginary $A_{\rm I}(\omega')$ parts as a function of $\omega' = \omega_d - \omega$ are plotted in (a) and (b), respectively, for two different values of Δ_d.

in the bistable regime.

Photon blockade and transparency.— To further discuss properties of the light field transmission when the photons are blockaded, we now study the response of the circuit QED system to the vacuum input field, which can be considered as a weak detecting field. According to the input-output theory [28], the output field with the Fourier components can be written as

$$a_{\rm out}(\omega) = A_{\rm in}(\omega') + \sqrt{2\gamma}A_f(\omega') + \left[\frac{2\gamma A_s + i\Omega}{\sqrt{2\gamma}}\right]\delta(\omega'), \tag{8}$$

with $\omega' = \omega + \omega_d$ and $A_f(\omega)$ given in Eq. (6). The physical meaning becomes clear if the vacuum input $a_{\rm in}(t)$ is assumed as a single-mode field $a_{\rm in}(t) = \xi\exp(-i\omega t)$ with the real parameter $\xi \ll |\Omega|$; that is, this weak detecting field does not change the statistical properties of the cavity field. In this case, ω' in Eq. (8) is changed to $\omega' = \omega_d - \omega$, and the terms $A_{\rm in}(\omega')$ and $\delta(\omega')$ in Eq. (8) denote the response of the system to the input vacuum field and the classical driving field, respectively. However, the term with $A^\dagger_{\rm in}(\omega')$ exhibits four-wave mixing with frequency $\omega - 2\omega_d$, which will be studied elsewhere.

The coefficient of $A_{\rm in}(\omega')$ for the output in Eq. (8) corresponds to that of $A_{\rm in}(\omega')$ in the expression $A_f(\omega')$ of Eq. (6) plus one. Thus in Figs. 4(a,b), the real $A_{\rm R}(\omega')$ and imaginary $A_{\rm I}(\omega')$ parts for the normalized coefficient of $A_{\rm in}(\omega)/\xi$ in Eq. (6) for $A_{\rm in}(t) = \xi\exp[-i(\omega - \omega_d)t]$ are plotted by using the same parameters as in Fig. 3(c), with $\Delta_d \approx 9.96$ MHz and $\Delta_d \approx 9.74$ MHz. These two values correspond to the minimum (photon blockade) and the maximum (classical case) of $g^{(2)}(0)$ in Fig. 3(c). As expected, we find that the response of the circuit QED system to the input field (or, say, weak-detecting field) has a Lorentzian shape, which is the same as the decay spectrum of the single photon, when the photon is blockaded. However, the weak detecting field shows transparency windows to the circuit QED system when the driving field is changed such that the cavity field is in the classical regime (or say the photon is not blockaded). Thus, we can control the photon (from blockade to transparency) by changing the applied classical field.

Conclusions.— In summary, we have studied the tunable transmission from the photon blockade to the photon transparency in superconducting circuit QED systems when the interaction between the qubit and the cavity field is in the dispersive regime. We analyze the effect of the driving field on the photon blockade and show the relation between the optical bistability and the photon blockade. We find that the photon blockade can be controlled by a classical driving field, that is, the photon blockade strongly depends on the properties of the driving field. We also find that the circuit QED system can be used to generate a four-wave mixing signal. All parameters in our numerical simulation are taken from experimentally available data. Therefore, our study should be experimentally realizable with current technology.

YXL is supported by the NSFC under Grant Nos. 10975080 and 61025022. AM is supported by the Polish Ministry of Science and Higher Education under Grant No. 2619/B/H03/2010/38. FN is partially supported by the LPS, NSA, ARO, NSF grant No. 0726909, JSPS-RFBR contract No. 09-02-92114, Grant-in-Aid for Scientific Research (S), MEXT Kakenhi on Quantum Cybernetics, and the JSPS via its FIRST program.

[1] J. Q. You and F. Nori, Phys. Today **58**, 42 (2005); Nature (London) **474**, 589 (2011).
[2] R. J. Schoelkopf and S. M. Girvin, Nature (London) **451**, 664 (2008).
[3] J. Bourassa, J. M. Gambetta, A. A. Abdumalikov, Jr., O. Astafiev, Y. Nakamura, and A. Blais, Phys. Rev. A **80**, 032109 (2009).
[4] T. Niemczyk, F. Deppe, H. Huebl, E. P. Menzel, F. Hocke, M. J. Schwarz, J. J. Garcia-Ripoll, D. Zueco, T. Hümmer, E. Solano, A. Marx, and R. Gross, Nature Phys. **6**, 772 (2010).
[5] P. Forn-Diaz, J. Lisenfeld, D. Marcos, J. J. Garcia-Ripoll, E. Solano, C. J. P. M. Harmans, and J. E. Mooij, Phys. Rev. Lett. **105**, 237001 (2010).
[6] S. Ashhab and F. Nori, Phys. Rev. A **81**, 042311 (2010).
[7] Y. X. Liu, C. P. Sun, and F. Nori, Phys. Rev. A **74**, 052321 (2006).
[8] C. M. Wilson, T. Duty, F. Persson, M. Sandberg, G. Johansson, and P. Delsing, Phys. Rev. Lett. **98**, 257003 (2007); C. M. Wilson, G. Johansson, T. Duty, F. Persson, M. Sandberg, and P. Delsing, Phys. Rev. B **81**, 024520 (2010).
[9] M. Hofheinz, E. M. Weig, M. Ansmann, R. C. Bialczak, E. Lucero, M. Neeley, A. D. O'Connell, H. Wang, J. M. Martinis, and A. N. Cleland, Nature (London) **454**, 310 (2008).
[10] J. M. Fink, M. Göppl, M. Baur, R. Bianchetti, P. J. Leek, A. Blais, and A. Wallraff, Nature (London) **454**, 315 (2008).
[11] L. S. Bishop, J. M. Chow, J. Koch, A. A. Houck, M. H. Devoret, E. Thuneberg, S. M. Girvin, and R. J. Schoelkopf, Nature Phys. **5**, 105 (2009).
[12] A. Imamoḡlu, H. Schmidt, G. Woods, and M. Deutsch, Phys Rev. Lett. **79**, 1467 (1997).
[13] K. M. Birnbaum, A. Boca, R. Miller, A. D. Boozer, T. E. Northup, and H. J. Kimble, Nature (London) **436**, 87 (2005).
[14] A. Faraon, I. Fushman, D. Englund, N. Stoltz, P. Petro, and J. Vuckovic, Nature Phys. **4**, 859 (2008).
[15] C. Lang, D. Bozyigit, C. Eichler, L. Steffen, J. M. Fink, A. A. Abdumalikov, M. Baur, S. Filipp, M. P. da Silva, A. Blais, and A. Wallraff, Phys. Rev. Lett. **106**, 243601 (2011).
[16] A. J. Hoffman, S. J. Srinivasan, S. Schmidt, L. Spietz, J. Aumentado, H. E. Treci, and A. A. Houck, Phys. Rev. Lett. **107**,

053602 (2011).

[17] E. Ginossar, L. S. Bishop, D. I. Schuster, and S. M. Girvin, Phys. Rev. A **82**, 022335 (2010).

[18] M. Boissonneault, J. M. Gambetta, and A. Blais, Phys. Rev. Lett. **105**, 100504 (2010).

[19] N. Lu, Phys. Rev. A **40**, 1707 (1989).

[20] H. Wang, M. Hofheinz, M. Ansmann, R. C. Bialczak, E. Lucero, M. Neeley, A. D. O'Connell, D. Sank, J. Wenner, A. N. Cleland, and J. M. Martinis, Phys. Rev. Lett. **101**, 240401 (2008).

[21] G. Compagno and F. Persico, Phys. Rev. A **22**, 2108 (1980).

[22] H. Ian, Y. X. Liu, and F. Nori, Phys. Rev. A **81**, 063823 (2010).

[23] Y. Nakamura, C. D. Chen, and J. S. Tsai, Phys. Rev. Lett. **79**, 2328 (1997).

[24] R. W. Boyd, *Nonlinear Optics* (Academic press, New York, 2008).

[25] C. W. Gardiner and M. J. Collett, Phys. Rev. A **83**, 063823 (2011).

[26] T. C. H. Liew and V. Savona, Phys. Rev. Lett. **104**, 183601 (2010).

[27] L. S. Bishop, E. Ginossar, and S. M. Girvin, Phys. Rev. Lett. **105**, 100505 (2010).

[28] D. F. Walls and G. J. Milburn, *Quantum Optics* (Springer, Berlin, 1994).